

Intergeneric Relations of the Angular - Toed Geckos of Circum Western Himalayas (Sauria: Gekkonidae)

Muhammad Sharif Khan*

Herpetological Laboratory, Rabwah 35460, Pakistan

Abstract.- Intergeneric relationships between angular-toed gekkonid genera of the western Himalayas are investigated. Cladistic analysis based on a set of external morphological characters indicates that *Altigekko* is most primitive genus and there exists a monophyletic generic lineage between genera *Altigekko*, *Sivaligekko*, *Cyrtopodion*, and *Indogekko* of circum western Himalayas.

Key words: Angular-toed geckos, western Himalayas, phylogeny.

INTRODUCTION

Geckos have always been a favorite group of animals with herpetologists because of their circum global distribution and frequent occurrence in the collection bags. Due to extensive work on geckos subfamilial (Underwood, 1954; Kluge, 1967) and intergeneric (Loveridge, 1947) relationships among most of the gekkotin groups are now quite established. However, there still remain some problematic groups: Afro-Malagasy (Bauer, 1990), Palearctic (Szczerbak and Golubev, 1996), and the western Himalayan geckos (Khan, 1993a,b, 2000, 2003b; Khan and Rösler, 1999). Recently *Cyrtopodion* geckos inhabiting Indus Valley, Pakistan, have been found to be monophyletic (Khan, 2003a). Similarly Macey *et al.* (2000) have established monophyly of Palearctic genera *Cyrtopodion* and *Mediodactylus*. Taxonomic problem of the angular-toed geckos of the western Himalayas (Anderson, 1872; Boulenger, 1890; Annandale, 1913; Smith, 1935; Minton, 1966; Mertens, 1969; Szczerbak and Golubev, 1986; Kluge, 1991, 1993, 2001; Rösler, 2000; Das, 1996; Schleich and Kästle 1998, 2002), has recently been solved (Khan, 2003b), by erecting three new genera *Indogekko*, *Sivaligekko* and *Altigekko*, solving a long standing taxonomic problem with gekkologists.

Present is the preliminary study to assay the inter-generic relationships among circum-

0030-9923/2009/0001-0029 \$ 8.00/0

Copyright 2009 Zoological Society of Pakistan.

Himalayan angular-toed new gekkotin genera *Cyrtopodion*, *Indogekko*, *Sivaligekko* and *Altigekko*.

MATERIALS AND METHODS

The morphological data used in present analysis were obtained from alcohol preserved specimens. Individual morphological character was assigned polarity using eublepharine geckos as distant outgroup. Sixteen character states were distinguished which were used to construct data matrix (Table I). Primitive and advanced states of the characters were determined by the criteria used and defined by Kluge (1967) and Szczerbak and Golubev (1996). A more detailed analysis awaits collection of fresh material from the western Himalayas, that may hardly differ from the conclusions drawn from present study.

Museum material used

Altigekko baturensis BMNH 1990.3 (MSK 007887), CAS 170529 (MSK 0769.87); *Altigekko stoliczkai* NMW 16756; *Altigekko yarkandensis* BMNH 72.3.22.4; *Cyrtopodion k. kachhense* BMNH 1990.7; *Cyrtopodion kachhense ingoldbyi* BMNH 1931.6.18.1-2, UF 71794, FMNH 235535; *Cyrtopodion caspius* SR 2546:16713-14; *Cyrtopodion fedtschenkoi* SR 1078:8837-8; *Cyrtopodion montiumsalsorum* BMNH 1904.11.19.1, USNM 257535, MSK 014.86; *Cyrtopodion turcmenicus* SR 961:8016-17; *Indogekko longipes voraginosus* CAS 130323;

* Address for correspondence: 306 N. Morton Ave MORTON PA, 190. typhlops99@hotmail.com

Indogekko l. longipes CAS 115944, SR 307:3267-68; *Indogekko fortmunroi* BMNH 1990.4 (MSK 0626.90); *Mediodactylus walli* BMNH 1910.7.12.1; *Mediodactylus chitralensis* BMNH 1946.8.23.19; *Siwaligekko dattanensis* MNHN 1979-7745; *Siwaligekko battalensis* BMNH 1990.2 (MSK 0764), SR 3046:20252 (MSK 0737.89), NMW 31720 (MSK 0762.89), CAS 170533 (MSK 0766.89), USNM 284135 (MSK 0767.89), FMNH 235534 (MSK 0765.89), MSK 0763.89.

Abbreviations used

BMNH, Natural History Museum, London; CAS, California Academy of Sciences, California, USA; FMNH, Field Museum of Natural History, Chicago, USA; MSK, Herp laboratory, 15/6 Darul Saddar North, Rabwah 35460, Pakistan (author's personal collection, now deposited in Natural History Museum, Government College University, Lahore, Pakistan); NMW, Naturhistorisches Museum Wien, Austria; SR, Institute of Zoology, Academy of Sciences, Kiev, Ukraine; UF, Florida State Museum, Gainesville, USA; USNM, National Museum of Natural History, Washington, D.C.

Definition and weight assignment to the character states

The following characters were considered. Primitive (primary) states are indicated by lower case letters, while advanced (derived) states by upper case letters, with number of asterisks indicating advance level of the derived state.

Character	Weight
1. Scale size (head and body)	
a, small (primitive)	1
A, large (advance)	2
2. Scale morphology	
b, homogenous (primitive)	1
B, Heterogeneous (advance)	2
3. Dorsal tubercle morphology	
C, Bead-like (primitive)	1
C, Flat keelless (primitive)	2
C*, Flat keeled (advance)	3
C**, Trihedral (advance)	4
4. Precloacal pores	
d, Precloacal pores only in male (primitive)	1
D, Precloacal pores in male and female (advance)	2
5. Femoral pores	
e, Femoral pores absent (primitive)	1
E, Femoral pores present (advance)	2
6. Basal phalanges	
f., Wide (primitive)	1
F, Narrow (advance)	2
7. Digital	
g, Poor (primitive)	1

	angulations	G, Acute (advance)	2
8.	Tail	h, Indistinct (primitive)	1
	segmentation	H, Distinct at base (advance)	2
		H ⁺ , Distinct throughout (advance)	3
9.	Dark band on the back of head	i, Present (primitive)	1
		I, Absent (advance)	2
10.	Subcaudals	j, Indistinct, in several rows (primitive)	1
		J, Distinct, in a single row (advance)	2
11.	Caudal tubercles	k, Small spiny (primitive)	1
		K, Large flat (advance)	2
		K*, Large, flat, keeled (advance)	3
		K**, Large, flat, keeled overlapping (advance)	4
12.	Limb size	l, Short, anterior reaching to eye (primitive)	1
		L, Long, anterior reaching naris and beyond (advance)	2
13.	Size of digits	m, Short (primitive)	1
		M, Long (advance)	2
14.	Snout slope	n, Gradual (primitive)	1
		N, Abrupt (advance)	2
15.	Cloacal sacs	o, Present (primitive)	1
		O, Absent (advance)	2
16.	Relative size of body and tail	p, Subequal (primitive)	1
		P, Tail longer (advance)	2

RESULTS AND DISCUSSION

The matrix generated on the basis of above characters is shown in Table I.

Despite counter interpretations of fossil records from Asia and Europe (Smith and Bryden, 1977; Estes, 1983), it is now strongly believed that southeast Asia played an important role in the evolution of major groups of reptiles: gekkonids (Kluge, 1967, 1987; Owen, 1983; Estes 1983; Grismer, 1988), agamids (Moody, 1980), varanids and scincids (Estes, 1983), and amphibians (Darlington, 1957; Blair, 1972; Savage, 1973; Khan, 1980). Geckos had extensive inter-island water-rafting activity in the Indian Ocean during the Cenozoic (Hahn *et al.*, 2004), as depicted in Figure 1. The ancestral aeluroscalabotens radiated in the north, diversified in Asian genus *Eublepharis*, and crossed into Central America via Beringia, as genus *Coleonyx* (Kluge, 1967; Grismer, 1988), while in west they reached Africa by early Tertiary as genus *Hemitheconyx* and *Holodactylus* (Grismer, 1988). However, genus *Cyrtodactylus* invaded extensively the oceanic islands by water-rafting, from island to island, across the Indian Ocean (Khan and Rösler, 1999). The oceanic cyrtodactylids reached

northeastern Gondwanaland by Miocene. Their xerophilic stock invaded drier north African sub Sahara, was probably ancestral to Indo-Palearctic




Fig. 1. Radiation and interrelationships between angular-toed gekkota of southeast Asia, and their paleo-zoogeography (modified from Estes, 1983; Fig. 6).

“cyrtopodion” geckos (Axelrod and Raven, 1974; Leviton and Anderson, 1984; Bobrov 2000; Khan 2002, 2003a). The African cyrtodactylids on the north-eastern Gondwanaland were rafted when it

broke off during Tertiary as “Indian Plate”. Carrying several tropical African Tertiary animals and plants as it moved across Indian Ocean (Blanford, 1876; Simpson, 1965; Estes, 1983; Karanth, 2003), which included geckos, agamids, skinks, varanids (Estes, 1983), and microhylid frogs (Savage, 1973). Briggs (2003), Prasad and Range (1995) and Range (1996) suggested that during late Cretaceous India remained connected to Eurasia, as well as to Madagascar and Seychelles plateau by a land bridge, as also suggested Upper Cretaceous invertebrates (Whatly and Bajpai, 2000).

The cyrtodactylids radiated across the subcontinent, speciating extensively on mainland and oceanic islands (Drevsky and Szczerback, 1997; Drevsky *et al.*, 1997; Schleich and Kastle, 1998; Bauer, 2002; Buer *et al.*, 2003; Pauwels *et al.*, 2004). Most probably *Siwaligecko* represent cyrtodactylids that inhabited oceanic islands falling in the way of drifting Indian plate, and were swept along and crushed between the plates (Buffetaut and Ingavat 1985; Jaeger *et al.*, 1989), are now represented by the subHimalayan mountainous crumple known as Siwaliks (Khan, 1979; Powel, 1979; Khan, 1980, 1993). The *Swaligecko* are widely distribution in Siwaliks from western to eastern Himalayas (Khan, 1993, 2003b, 2005, 2006). The geckos on the northern part of the plate were carried along the Himalayan massif to high altitudes and now are represented by genus *Altigecko* (Khan, 2004, 2006). Widely distributed peculiar sandstone geckos of genus *Indogecko* (Khan, 2006), have strong affinities with cyrtopodion geckos that invaded circum Indus region (Khan, 2003).

Table I rates the proximity of these genera to the ancestral southeast Asian geckos represented by *Eublepharis macularious* in the analysis. *Altigecko* are closest to ancestral form while *Indogecko* and *Cyrtopodion* occupy farthest position in the spectrum.

REFERENCES

- ANDERSON, J., 1872. On some Persian, Himalayan and other reptiles. *Proc. zool. Soc. London*, **1872**: 417-498.
- ANNANDALE, N. 1913. The Indian geckoes of the genus *Gymnodactylus*. *Rec. Indian Mus., Calcutta*, **9**: 309-362.

AXELROD, C.I. AND RAVEN, P.H., 1974. Late Cretaceous and Tertiary vegetation history of Africa. In: *Biogeography and ecology of southern Africa* (ed.

Table I.- Morphometric analysis of angular-toed gekkonid genera of circum western Himalayas.

Genera		Characters																Total weight
		a-A	b-B	-C-C*-C	d-D	e-E	f-F	g-G	h-H-H*	i-I	j-J	k-K*-K*	l-L	m-M	n-N	o-O	p-P	
<i>Eublepharis</i> (Ncestral)	States	a	b	c	d	e	f	g	h	i	j	k	l	m	n	o	p	16
	Weight	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	
<i>Altigekko</i>	States	a	b	c	d	e	f	g	H	i	j	k	l	m	n	o	p	18
	Weight	1	1	1	1	1	1	1	2	2	1	1	1	1	1	1	1	
<i>Siwaligekko</i>	States	a	b	c	d	e	f	g	h	i	j	K	l	m	N	o	p	19
	Weight	1	1	1	1	1	1	1	1	1	2	1	2	1	2	1	1	
<i>Indogekko</i>	States	A	B	C-C*	D	E	F	G	H*	i	J	K*	L	M	N	O	P	38
	Weight	2	2	2+3	2	2	2	2	3	1	2	3	2	2	2	2	2	
<i>Cyrtopodion</i>	States	A	B	C**	D	E	F	G	H	I	J	K	L	M	N	O	p	35
	Weight	2	2	4	1	2	2	2	3	2	2	4	2	2	2	2	1	

- M.J.H. Weaver) W Junk Publishers, The Hague, pp. 77-131.
- BAUER, A.M., 1990. Phylogeny and biogeography of the geckos of southern Africa and the islands of the western Indian Ocean: A preliminary analysis. In: *Vertebrates in the tropics* (eds. G. Peter and R. Hutterer) Museum Alexander Koenig, Bonn, pp. 275-284.
- BAUER, A.M., 2002. Two new species of *Cyrtodactylus* (Squamata: Gekkonidae) from Myanmar. *Proc. Calif. Acad. Sci.*, **53**: 73-86.
- BAUER, A.M., SUMOTHA, M. AND PAUWELS, O.G., 2003. Two new species of *Cyrtodactylus* (Reptilia: Squamata: Gekkonidae) from Thailand. *Zootaxa*, **376**: 1-18.
- BLAIR, W.F., 1972. *Evolution in the genus Bufo*. University of Texas Press, Austin and London.
- BLANFORD, W.T., 1876. The African element in the fauna of India: A criticism of Mr. Wallace's views as expressed in the 'Geographical Distribution of Animals'. *Annls Mag. Nat. Hist.*, **18** (4th ser.): 277-294.
- BOBROV, V.V., 2000. The limit between Palearctic and Afrotropical faunistic kingdoms (according to distribution of lizards (Reptilia, Sauria)). *Biol. Bull.*, **27**: 151-163.
- BOULENGER, G.A., 1890. *Fauna of British India, including Ceylon and Burma. Reptilia and Batrachia*. London.
- BRIGGS, J.C., 2003. The biographic and tectonic history of India. *J. Biogeogr.*, **30**: 381-388.
- BUFFETAUT, E. AND INGAVAT, R., 1985. The Mesozoic vertebrates of Thailand. *Sci. Am.*, **251**: 64-71.
- DAS, I., 1996. *Biogeography of the reptiles of South Asia*. Krieger Publishing Company, Malabar, Florida.
- DAREVSKY, I.S., HELFEBERGER, N., ORLOV, N. AND SHAH, K., 1997. Two new species of genus *Gonydactylus* (Sauria: Gekkonidae) from Eastern Nepal. *Russian J. Herpetol.*, **4**: 89-93.
- DAREVSKY, I.S. AND SZCZERBAK, N.N., 1997. A new gecko of the genus *Gonydactylus* (Sauria: Gekkonidae) with a key to the species from Vietnam. *Asiat. Herpetol. Res.*, **7**: 19-22.
- DARLINGTON, P.J. Jr. 1957. *Zoogeography: the geographical distribution of animals*. John Wiley, N.Y.
- ESTES, R., 1983. The fossil record and early distribution of lizards. In: *Advances in herpetology and evolutionary biology* (eds. A.J.G. Rhodin and K. Miyata). Museum of Comparative Zoology, Cambridge, Mass, pp. 365-398.
- GRISMER, L.L., 1988. The phylogeny, taxonomy classification, and biography of eublepharid geckos (Reptilia: Squamata). In: *Phylogenetic relationships of the lizard families* (eds. R.D. Estes and G.K. Pregill). Stanford University Press, Stanford, pp. 369-469.
- HAHN, D., ZHOU, K. AND BAUER, A. M., 2004. Phylogenetic relationships among gekkotan lizards inferred from *Cmos* nuclear DNA sequences and a new classification of the Gekkota. *Biol. J. Linn. Soc.*, **83**: 353-368.
- JAEGER, J.J., COURTILLOT, V. AND TAPPONNIER, P., 1989. Paleontological view of the ages of the Deccan Traps the Cretaceous/Tertiary boundary, and the India-

- Asia collision. *Geology*, **17**: 316-319.
- KARANTH, K.P., 2003. Evolution of disjunct distributions among wet-zone species of the Indian subcontinent: Testing various hypotheses using a phylogenetic approach. *Curr. Sci.*, **85**: 101-108.
- KHAN, M. S., 1979. On a collection of amphibians from northern Punjab and Azad Kashmir, with ecological notes. *Biologia (Lahore)*, **25**: 37-50.
- KHAN, M.S., 1980. Affinities and zoogeography of herpetiles of Pakistan. *Biologia (Lahore)*, **26**: 113-171.
- KHAN, M.S., 1988. A new cyrtodactylid gecko from north-western Punjab, Pakistan. *J. Herpetol.*, **22**: 241-243.
- KHAN, M.S., 1993a. A new angular-toed gecko from Pakistan, with remarks on the taxonomy and a key to the species belonging to genus *Cyrtodactylus* (Reptilia:Sauria: Gekkonida). *Pakistan J. Zool.*, **25**: 67-73.
- KHAN, M.S., 1993b. A new sandstone gecko from Fort Munro, Dera Ghazi Khan District, Punjab, Pakistan. *Pakistan J. Zool.*, **25**: 217-221.
- KHAN, M.S., 2000. Redescription and generic redesignation of *Gymnodactylus stoliczkai* Steindachner, 1869. *Pakistan J. Zool.*, **32**: 157-163.
- KHAN, M.S., 2001. Taxonomic notes on angulat-toed gekkota of Pakistan, with description of a new species of genus *Cyrtopodion*. *Pakistan J. Zool.*, **33**: 13-24.
- KHAN, M.S., 2003a. Notes on circum Indus geckos of genus *Cyrtopodion* (Squamata: Gekkonidae). *Gekkota, Quedlinburg*, **4**: 43-59.
- KHAN, M.S., 2003b. Questions of generic designation of angular-toed geckos of Pakistan with descriptions of three new genera (Reptilia: Gekkonidae). *J. nat. Hist. Wildl. (Karachi)*, **2**: 1-9.
- KHAN, M.S., 2004. Notes on high altitude geckos of the genus *Altigecko* in northeastern Pakistan. *Bull. Chicago Herpetol. Soc.*, **39**: 221-224.
- KHAN, M.S., 2005. Notes on geckos of the genus *Siwaligecko* Khan, 2003 (Reptilia: Gekkonidae) in Pakistan. *Bull. Chicago Herpetol. Soc.*, **40**: 54-57.
- KHAN, M.S., 2006. *Amphibians and reptiles of Pakistan*. Krieger Publishing Company, Malabar, Florida.
- KHAN, M.S., 2006. Biology and distribution of geckos of genus *Indogecko* Khan, 2003 (Sauria: Gekkonidae). *Russian J. Herpetol.*, **15**: 87-92.
- KHAN, M.S. AND RASHIDA, T., 1990b. A new gecko of the genus *Tenuidactylus* from northeastern Punjab, Pakistan, and southwestern Azad Kashmir. *Herpetologica*, **46**: 142-148.
- KHAN, M.S. AND RÖSLER, H., 1999. Redescription and generic reallocation of Ladakhian gecko *Gymnodactylus stoliczkai* Steindachner, 1869. *Asiatic Herpetol. Res.*, **8**: 60-68.
- KLUGE, A.G., 1967. Higher taxonomic categories of gekkonid lizards and their evolution. *Bull. Am. Mus. nat. Hist.*, **135**: 1-60.
- KLUGE, A.G., 1987. Cladistic relationships in the Gekkonoidea (Squamata, Sauria). *Misc. Publ. Mus. Zool., Univ. Michigan*, **173**: 1-54.
- KLUGE, A.G., 1991. Checklist of gekkonoid lizards. *Smithsonian Herpetol. Inf. Serv.*, **85**: 1-35.
- KLUGE, A.G., 1993. *Gekkonid lizard taxonomy*. International Gecko Society, San Diego, California.
- KLUGE, A.G. 2001. Gekkotan lizard morphology. *Hamadryad*, **26**: 1-209.
- LEVITON, A. E. AND ANDERSON, S.C., 1984. Description of a new species of *Cyrtodactylus* from Afghanistan with remarks on the status of *Gymnodactylus longipes* and *Cyrtodactylus fedtschenkoi*. *J. Herpetol.*, **18**: 270-276.
- LOVERIDGE, A., 1947. Revision of the African lizards of the family Gekkonidae. *Bull. Mus. Comp. Zool.*, **98**: 1-469, 7 plates.
- MACEY, J.R., ANNANJEVA, N.B., WANG, Y. AND PAPENFUSS, T.J., 2000. Phylogenetic relationships among Asian gekkonid lizards formerly of the genus *Cyrtodactylus* based on cladistic analyses of allozymic data: monophyly of *Cyrtodactylus* and *Mediodactylus*. *J. Herpetol.*, **34**: 258-265.
- MERTENS, R., 1969. Die Amphibien und Reptilien West-Pakistans. *Stuttg. Beitr. Naturk.*, **197**: 1-96.
- MINTON, S.A., 1966. A contribution to the herpetology of West Pakistan. *Bull. Am. Mus. nat. Hist.*, **134**: 31-184.
- MOODY, S.M., 1980. *Phylogenetic and historical biogeographical relationships of the genera in the family Agamidae (Reptilia: Lacertilia)*. Doctoral Dissertation, University of Michigan, Ann Arbor, pp. 373.
- OWEN, H.G., 1983. *Atlas of continental displacement 200 million to the present*. Cambridge University Press, Cambridge.
- PAUWELS, O.G., BAUER, A., SUMONTHA, M. AND CHANHOM, L., 2004. *Cyrtodactylus thirakhupti* (Squamata: Gekkonidae), a new cave-dwelling gecko from southern Thailand. *Zootaxa*, **772**: 1-11.
- POWEL, C.M., 1979. A speculative tectonic history of Pakistan and surroundings: some constraints from the Indian Ocean. In: *Geodynamics of Pakistan*. Geological Survey of Pakistan, Sp. Memo., pp. 5-25.
- PRASAD, G.V. AND RANGE, J.C., 1995. Amphibians and squamates from the Maastrichtian of Naskal, India. *Cretaceous Res.*, **16**: 95-107.
- RAGE, J.C., 1996. Le peuplement animal de Madagascar: une composante venue de Laurasia est-elle envisageable? In: *Biogéographie de Madagascar* (edt. W.R. Lourenco). OSTROM, Paris, pp. 27-35.
- RÖSLER, H., 2000. Kommentierte Liste der rezent, subrezent und Fossil bekannten Geckotaxa (Reptilia: Gekkonomorpha). *Gekkota*, **2**: 28-153.
- SAVAGE, J.M., 1973. The geographic distribution of frogs:

- patterns and predictions. In: *Evolutionary biology of the anurans, contemporary research and major problems* (ed. J.L. Vial), Chapter 13. University Missouri Press, Columbia, pp. 351-445.
- SCHLEICH, H.H. AND KÄSTLE, W., 1998. Description of *Gonydactylus nepalensis* sp. nov. from the inner Terai of far West Nepal (Reptilia: Sauria: Gekkonidae). *Veroffentl. Fuhlrott-Mus.*, **4**: 269-280.
- SCHLEICH, H. H. AND KÄSTLE, W., 2002. *Amphibians and reptiles of Nepal*. Koeltz Scientific Books, Koenigstein, Germany, pp. 1-1200.
- SIMPSON, G.G., 1965. *The geography of evolution: collected essays*. Chilton Books, Philadelphia.
- SMITH, M.A. 1935. *The Fauna of British India, including Ceylon and Burma. Reptilia and amphibia*. Vol. II: *Sauria*. Taylor and Francis Ltd. London.
- SMITH, A. AND BRYDEN, J., 1977. *Mesozoic and cenozoic paleocontinental maps*. Cambridge University Press, pp. 70.
- SZCZERBAK, N.N. AND GOLUBEV, M.L., 1984. On generic assignment and intergeneric structure of the Palearctic *Cyrtodactylus* lizards. *Vest. Zool.*, **2**: 50-56 (in Russian).
- SZCZERBAK, N.N. AND GOLUBEV, M. L., 1986. Geckos of USSR fauna and adjoining countries. *Sci. Acad. Ukr. SSR Zool. Inst.*, **1986**: 1-232 (in Russian).
- SZCZERBAK, N.N. AND GOLUBEV, M. L., 1996. *Gecko fauna of the USSR and contiguous regions*. Translation. Soc. Study Amphib. Rept. pp. 1-233.
- UNDERWOOD, G. 1954. On the classification and evolution of geckos. *Proc. zool. Soc. London*, **124**: 469-492.
- WHATLEY, R.C. AND BAIPAI, S., 2000. Zoogeographical relationships of the Upper Cretaceous non marine Ostracoda of India. *Curr. Sci.*, **79**: 694-696.

(Received 14 July 2008, revised 15 October 2008)